AU InforMed

Volume 10 Number 3 (Issue 254)

Monday, June 4, 2012

Guest Editor: Katherine (Katie) Teel, Pharm.D.

Key Inforbits

• New drugs approved by the FDA in 2011

• New FDA-Approved Dosage Forms in 2011

New Biologicals approved in 2011

'The Last Dose'

2011 New FDA Drug Approvals.

| Arcapta Neohaler (indacaterol) Brilinta (iticagrelor) AstraZeneca (iticagrelor) Reversible P2Y ₁₂ antagonist (iticagrelor) AstraZeneca (iticagrelor) AstraZeneca (iticagrelor) AstraZeneca (iticagrelor) AstraZeneca (iticagrelor) Multikinase inhibitor Medullary thyroid cancer (vandetanib) Caoper Medullary thyroid cancer Reversible P2Y ₁₂ antagonist (vandetanib) Caoper Medullary thyroid cancer Reversible P2Y ₁₂ antagonist (vandetanib) Caoper Medullary thyroid cancer Reversible via the department of thrombotic events Routine use of antidiarrhea is recommended with this agent. | Drug | Manufacturer | Mechanism | Indications | Other Information |
|--|-------------------|-------------------|---|-----------------------|-----------------------------------|
| Reversible P2Y12 antagonist (indacaterol) | _ | Novartis | | COPD | |
| Brilinta (ticagrelor) | | | (LABA) | | related deaths, like other LABAs. |
| Caprelsa (vandetanib) | | | | | |
| Caprelsa (vandetanib) | | AstraZeneca | Reversible P2Y ₁₂ antagonist | | _ |
| (vandetanib) Forest PDE-IV inhibitor COPD Weight loss within 6 months and diarrhea within 4 weeks are major ADRs. DaTscan (ioflupane I 123) GE Healthcare Binds reversibly to the dopamine transporter Brain diagnostic imaging Adjunct in the differential diagnosis of essential tremor from Parkinsonian syndromes. Difficid (fidaxomicin) Optimer Pharm. Bactericidal macrolide antibiotic C. difficile associated diarrhea Clinical trials found fidaxomicin non-inferior to vancomycin for C. difficile. Edarbi (azilsartan) Takeda Angiotensin-II Receptor Blocker Hypertension BBW against use in pregnancy, same as other ARBs. Edurant (rilpivirine) Tibotec Non-nucleoside Reverse Transcriptase Inhibitor HIV Must be taken with a 400-calorie meal for best absorption. Ferriprox (deferiprone) ApoPharma Genetic Therapies Selective Bradykinin B2 Receptor Antagonist Hereditary angioedema Administered as subcutaneous injection; Decreases efficacy of ACEIs. Gadavist (gadobutrol) Bayer Gadolinium-containing contrast agent Diagnostic Imaging Stemic fibrosis. BBW for increased risk of nephrogenic systemic fibrosis. Horizant (gabapentin encarabil) GSK Prodrug of gabapentin encarabil) Restless Leg Syndrome Once daily dosing; recommended with food at 5:00 p. | · · · · · · | | | | • • |
| Daliresp (roflumilast) Forest PDE-IV inhibitor COPD Weight loss within 6 months and diarrhea within 4 weeks are major ADRs. | _ | AstraZeneca | Multikinase inhibitor | • • | |
| Columnia Columnia | | | | | · · |
| dopamine transporter imaging essential tremor from Parkinsonian syndromes. | _ | Forest | PDE-IV inhibitor | COPD | |
| Dificid (fidaxomicin) | DaTscan | GE Healthcare | • | <u> </u> | |
| (fidaxomicin) Takeda Angiotensin-II Receptor Blocker Blocker Tibotec Non-nucleoside Reverse (rilpivirine) Ferriprox (deferiprone) ApoPharma (icatibant) Genetic Therapies Gadoutrol) Horizant (gabapentin enacarbil) Incivek (telaprevir) Jakafi Takeda Angiotensin-II Receptor Blocker Angiotensin-II Receptor Blocker HIV Must be taken with a 400-calorie meal for best absorption. BBW for agranulocytosis; monitor absolute neutrophil count. Hereditary angioedema Chronic Hepatitis C Substrate and strong inhibitor of CYP3A4. Myelofibrosis Dose adjustments made based upon | (ioflupane I 123) | | | | syndromes. |
| Takeda Angiotensin-II Receptor Blocker Hypertension BBW against use in pregnancy, same as other ARBs. | Dificid | Optimer Pharm. | | | |
| Blocker Other ARBs. | (fidaxomicin) | | | | • " |
| Tibotec Non-nucleoside Reverse Transcriptase Inhibitor Transcriptase Inhibitor Transcriptase Inhibitor Transcriptase Inhibitor Transcriptase Inhibitor Serine protease inhibitor Transcriptase Inhibitor Trans | | Takeda | | Hypertension | |
| Transcriptase Inhibitor BBW for agranulocytosis; monitor ApoPharma Oral iron chelating agent Iron overload BBW for agranulocytosis; monitor absolute neutrophil count. | (azilsartan) | | | | |
| Ferriprox (deferiprone) | | Tibotec | | HIV | |
| Administered as subcutaneous injection; Decreases efficacy of ACEIs. | | | * | | * |
| (icatibant) Genetic Therapies Receptor Antagonist Decreases efficacy of ACEIs. Gadavist (gadobutrol) Bayer Gadolinium-containing contrast agent Diagnostic Imaging Systemic fibrosis. Horizant (gabapentin (gabapentin enacarbil) Prodrug of gabapentin Prodrug of gabapentin (gabapentin enacarbil) Chronic Hepatitis C Substrate and strong inhibitor of CYP3A4. Jakafi Incyte Kinase inhibitor Myelofibrosis Dose adjustments made based upon | _ | ApoPharma | Oral iron chelating agent | Iron overload | • |
| Gadavist (gadobutrol) Horizant (gabapentin enacarbil) Incivek (telaprevir) Jakafi Bayer Gadolinium-containing contrast agent Gadolinium-containing contrast agent Bayer Gadolinium-containing contrast agent Bayer Gadolinium-containing contrast agent Bayer Gadolinium-containing systemic fibrosis. Restless Leg Syndrome Once daily dosing; recommended with food at 5:00 p.m. Chronic Hepatitis C Substrate and strong inhibitor of CYP3A4. Myelofibrosis Dose adjustments made based upon | Firazyr | | - | Hereditary angioedema | 3 |
| (gadobutrol)contrast agentsystemic fibrosis.Horizant (gabapentin enacarbil)GSKProdrug of gabapentin (gabapentin enacarbil)Restless Leg Syndrome food at 5:00 p.m.Once daily dosing; recommended with food at 5:00 p.m.Incivek (telaprevir)Vertex Pharm.Serine protease inhibitorChronic Hepatitis C CYP3A4.Substrate and strong inhibitor of CYP3A4.JakafiIncyteKinase inhibitorMyelofibrosisDose adjustments made based upon | (icatibant) | Genetic Therapies | 1 0 | | · · |
| (gabapentin enacarbil) Incivek Vertex Pharm. Serine protease inhibitor Chronic Hepatitis C Substrate and strong inhibitor of CYP3A4. Jakafi Incyte Kinase inhibitor Myelofibrosis Dose adjustments made based upon | | Bayer | C | Diagnostic Imaging | |
| enacarbil) Incivek (telaprevir) Jakafi Incyte Kinase inhibitor Myelofibrosis Vertex Pharm. Serine protease inhibitor Chronic Hepatitis C Substrate and strong inhibitor of CYP3A4. Myelofibrosis Dose adjustments made based upon | Horizant | GSK | Prodrug of gabapentin | Restless Leg Syndrome | |
| Incivek (telaprevir)Vertex Pharm.Serine protease inhibitorChronic Hepatitis CSubstrate and strong inhibitor of CYP3A4.JakafiIncyteKinase inhibitorMyelofibrosisDose adjustments made based upon | (gabapentin | | | | food at 5:00 p.m. |
| (telaprevir) CYP3A4. Jakafi Incyte Kinase inhibitor Myelofibrosis Dose adjustments made based upon | enacarbil) | | | | |
| Jakafi Incyte Kinase inhibitor Myelofibrosis Dose adjustments made based upon | | Vertex Pharm. | Serine protease inhibitor | Chronic Hepatitis C | <u>~</u> |
| Valuati V | | | | | |
| (ruxolitinib) degree of thrombocytopenia. | | Incyte | Kinase inhibitor | Myelofibrosis | |
| | (ruxolitinib) | | | | degree of thrombocytopenia. |

| Drug | Manufacturer | Mechanism | Indications | Other Information |
|--------------------------------|---------------------------|--|--|---|
| Natroba (spinosad) | ParaPRO/Pernix | Pediculocidal and ovicidal | Head lice | Topical shampoo approved for children >6 months. |
| Onfi (clobazam) | Lundbeck | Benzodiazepine | Lennox-Gastaut & Epilepsy (adjunct) | Weight-based dosing for LGS and pediatric epilepsy. |
| Potiga (ezogabine) | GSK | Neuronal K+ channel opener | Partial-onset seizures (adjunct) | CNS ADRs of dizziness, somnolence, and fatigue are common. |
| Tradjenta (linagliptin) | Boehringer Ingelheim | DPP-4 Inhibitor | T2DM | Same MOA as Januvia and Onglyza. |
| Victrelis (boceprevir) | Merck | Protease Inhibitor | Chronic Hepatitis C | Substrate and strong inhibitor of CYP3A4. |
| Viibryd (vilazodone) | Trovis | SSRI and 5HT _{1A} partial agonist | Depression | Decreased incidence of sexual dysfunction vs. other SSRIs in trials. |
| Xalkori (crizotinib) | Pfizer | Tyrosine kinase receptor inhibitor | Non-small cell lung cancer | Monitor for hematologic toxicity, LFT elevation >5X ULN, QT prolongation. |
| Xarelto (rivaroxaban) | Janssen | Factor Xa Inhibitor | Atrial fibrillation & post-op thromboprophylaxis | BBW for increased stroke risk after d/c of therapy and for hematoma occurrence with neuraxial anesthesia. |
| Zelboraf (vemurafenib) | Genentech | BRAF Kinase Inhibitor | Advanced Melanoma | Used in patients with unresectable or metastatic melanoma with a BRAF ^{V600E} mutation. |
| Zytiga (abiraterone) | Centocor Ortho Biotech | Irreversible CYP17 inhibitor | Advanced Prostate Cancer | Adjust dose based upon degree of hepatotoxicity after initiation. |

New Formulations:

• In 2011, several new dosage forms and combination products of previously approved drugs were also introduced. A few highlights from the new preparations list include:

| Drug | Dosage Form | Manufacturer | Indication |
|---------------------------------------|-------------------|---|---|
| Abstral | Sublingual tablet | ProStrakan | Breakthrough pain in opioid-tolerant cancer patients. |
| (fentanyl) | | | |
| AndroGel | 1.62% Topical Gel | 1.62% Topical Gel Abbott Increased strength, therapy fo | |
| (testosterone) | | | |
| Complera | Tablet | Gilead Sciences | Once daily, three-drug therapy for treatment-naïve |
| (emtricitabine/rilpivirine/tenofovir) | | | HIV infection. |
| Forfivo XL | 450 mg tablet | Intelgenx Corp. | Increased strength, extended-release therapy for |
| (bupropion) | | | depression. |
| Juvisync | Tablet | Merck | Combination tablet for type 2 diabetics with |
| (sitagliptin/simvastatin) | | | concomitant dyslipidemia. |
| Oxecta | Tablet | King | Immediate-release tablet formulated to reduce abuse |
| (oxycodone) | | Pharmaceuticals | potential. |
| Zutripro | Suspension | Cypress | Combination antitussive, antihistamine, and |
| (hydrocodone/chlorpheniramine/ | | Pharmaceuticals | decongestant for relief of allergy and cold symptoms. |
| pseudoephedrine) | | | |

New Biologicals:

| Name | Manufacturer | Mechanism | Indications | Other Information |
|--|------------------------------|---|---|--|
| Adcetris (brentuximab vedotin) | Seattle Genetics | Antibody drug conjugate directed at CD30 | Hodgkin's Lymphoma and systemic anaplastic large cell lymphoma after prior treatment failure | BBW for increased incidence of Progressive multifocal encephalopathy. |
| Anascorp (centruroides F(ab)2) | Rare Disease Therapeutics | Venom-specific F(ab)2 fragments of IgG that neutralize venom toxins | Antivenom for scorpion stings | Given IV over ten minutes as soon after scorpion sting as possible. |
| Benlysta (belimumab) | Human Genome Sciences/GSK | B-lymphocyte stimulator- specific inhibitor | Systemic lupus erythematosus | Adjunct to standard SLE therapy in autoantibody-positive SLE. |
| Corifact (factor XIII) | CSL Behring | Endogenous plasma glycoprotein that is converted to factor XIIIa in the presence of Ca ⁺² | Congenital factor XIII deficiency | Dose is adjusted based on factor XIII trough levels which are tested every 4 weeks. |
| Erwinaze (asparaginase Erwinia chrysanthemi) | EUSA Pharma | Antineoplastic agent that reduces levels of asparagine | Acute Lymphoblastic Leukemia in patients with hypersensitivity to <i>E.coli</i> - derived asparaginase | Hypersensitivity reactions occur in >10% of patients. |
| Eylea (aflibercept) | Regeneron Pharm. | Vascular endothelial growth factor inhibitor | Neovascular age-related macular degeneration | For intravitreal injection only; associated with endophthalmitis and retinal detachment. |
| Fluzone Intradermal (influenza vaccine) | Sanofi Pasteur | Induces specific antibody production versus influenza | Active immunity to vaccine-containing strains of influenza virus | Alternative to IM influenza vaccination in 18-64 yo. |
| Nulojix (belatacept) | BMS | Selective T-Cell Costimulation Blocker | Prophylaxis of organ rejection in kidney transplant recipients | Used concomitantly with basiliximab, mycophenolate, and corticosteroids in Epstein-Barr Virus seropositive patients. |
| Yervoy (ipilimumab) | BMS | Cytotoxic T-lymphocyte antigen-4 blocking antibody | Unresectable or metastatic melanoma | Can cause severe immune-mediated ADRs including dermatologic, endocrine, GI, hepatic, and ophthalmic toxicities. |

References:

- 2011 Novel New Drugs. January 2012. U.S. Department of Health and Human Services. U.S. Food and Drug Administration. Center for Drug Evaluation and Research. Silver Spring, MD. [last updated 2012 Jan; accessed 2012 Mar 14]. Available at:
 - $\frac{http://www.fda.gov/downloads/Drugs/DevelopmentApprovalProcess/DrugInnovation/UCM29366}{3.pdf}$
- 2011 Biological License Application Approvals. U.S. Department of Health and Human Services. U.S. Food and Drug Administration. Silver Spring, MD. [last updated 2012 Jan; accessed 2012 Mar 14]. Available at:
 - http://www.fda.gov/BiologicsBloodVaccines/DevelopmentApprovalProcess/BiologicalApprovalsby Year/ucm242933.htm
- New Drugs Approved by the FDA in 2011. Pharmacists Letter. Stockton, CA. Therapeutic Research Center. 2011. Available at:
 - http://pharmacistsletter.therapeuticresearch.com/pl/NewDrugs.aspx?cs=&s=PL&pt=20&yr=2011
- Hussar DA. New therapeutic agents marketed in the United States in 2011. Pharmacist Activist. 2011
 Dec;6(12):2-3. Available at: www.pharmacistactivist.com



The last "dose" ...

"Lost wealth may be replaced by industry, lost knowledge by study, lost health by temperance or medicine; but lost time is gone forever."



-Samuel Smiles, Scottish author and reformer (1812-1904)



"Ah, summer, what power you have to make us suffer and like it."

~Russell Baker (American writer, 1925 -)

An electronic bulletin of drug and health-related news highlights, a service of ...

Auburn University, Harrison School of Pharmacy, Drug Information Center

• Phone 334-844-4400 • Fax 334-844-8366 • http://www.pharmacy.auburn.edw/dilrc/dilrc.htm

Bernie R. Olin, Pharm.D., Director

Archived issues are available at: http://pharmacy.auburn.edu/dilrc/au informed.htm